

<https://helda.helsinki.fi>

---

## Habitat suitability modeling of Asian Elephant *Elephas maximus* (Mammalia: Proboscidea: Elephantidae) in Parsa National Park, Nepal and its buffer zone

Sharma, Puja

2019-10-26

---

Sharma , P , Adhikari , H , Tripathi , S , Ram , A & Bhattarai , R 2019 , ' Habitat suitability modeling of Asian Elephant *Elephas maximus* (Mammalia: Proboscidea: Elephantidae) in Parsa National Park, Nepal and its buffer zone ' , Journal of threatened taxa , vol. 11 , no. 13 pö , 10.11609/jott.4467.11.13.14643-14654 , pp. 14643 14654 . <https://doi.org/10.11609/jott.4467.11.13.14643-14654> .

---

<http://hdl.handle.net/10138/310445>

<https://doi.org/10.11609/jott.4467.11.13.14643-14654>

---

cc\_by

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

# Journal of Threatened Taxa

Building evidence for conservation globally

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

## COMMUNICATION

### HABITAT SUITABILITY MODELING OF ASIAN ELEPHANT *Elephas maximus* (MAMMALIA: PROBOSCIDEA: ELEPHANTIDAE) IN PARSA NATIONAL PARK, NEPAL AND ITS BUFFER ZONE

Puja Sharma, Hari Adhikari, Shankar Tripathi, Ashok Kumar Ram & Rajeev Bhattarai

26 October 2019 | Vol. 11 | No. 13 | Pages: 14643–14654

DOI: 10.11609/jott.4467.11.13.14643-14654



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact [ravi@threatenedtaxa.org](mailto:ravi@threatenedtaxa.org)

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

## Partner



صندوق محمد بن زايد  
للمحافظة على  
الكائنات الحية

The Mohamed bin Zayed  
SPECIES CONSERVATION FUND

## Member



## Publisher & Host







# HABITAT SUITABILITY MODELING OF ASIAN ELEPHANT *Elephas maximus* (MAMMALIA: PROBOSCIDEA: ELEPHANTIDAE) IN PARSA NATIONAL PARK, NEPAL AND ITS BUFFER ZONE

Puja Sharma<sup>1</sup> , Hari Adhikari<sup>2</sup> , Shankar Tripathi<sup>3</sup> , Ashok Kumar Ram<sup>4</sup>  &  
Rajeev Bhattarai<sup>5</sup> 

ISSN 0974-7907 (Online)  
ISSN 0974-7893 (Print)

PLATINUM  
OPEN ACCESS



<sup>1,3</sup> Faculty of Forestry, Agriculture and Forestry University, Hetauda, Nepal.

<sup>2</sup> Earth Change Observation Laboratory, Department of Geosciences and Geography, University of Helsinki, Finland.

<sup>2</sup> Institute for Atmospheric and Earth System Research, Faculty of Science, University of Helsinki, Finland.

<sup>4</sup> Conservation Officer, Department of National Parks and Wildlife Conservation Nepal.

<sup>5</sup> Tribhuvan University, Institute of Forestry, Pokhara, Nepal.

<sup>1</sup>oujasharma@gmail.com (corresponding author), <sup>2</sup>hari.adhikari@helsinki.fi, <sup>3</sup>stripathi@afu.edu.np,

<sup>4</sup>ashokrink11@gmail.com, <sup>5</sup>bhattarairajeev@gmail.com

**Abstract:** Asian Elephants *Elephas maximus* in Nepal are known to have habitats and movement corridors in Parsa National Park (PNP) and its buffer zone (BZ), located east of Chitwan National Park. A study was conducted in this area to assess the suitability of PNP and BZ as elephant use areas, and to determine factors relevant to the presence of elephants in PNP. Field measurements were carried out in 67 plots for vegetation analysis. Boosted Regression Tree (BRT) analysis was used to examine the relationship of habitat suitability and variables including topography (slope, aspect, altitude), climate (precipitation, temperature), habitat preference, ground cover and crown cover. The results indicate that elephant habitat suitability is mainly determined by the dominant plant species, temperature, altitude, habitat preference and precipitation. Slope, ground cover, crown cover and substrate have lesser effects. Elephants were recorded up to 400m in the northeast and southeast aspects of the study area. Most suitable habitats were low slope forest dominated by *Acacia catechu* and *Myrsine semicarpata* that received 300mm annual precipitation. The model emphasizes environmental suitability, and contributes to knowledge for conservation of elephants in PNP and BZ by delineating sites that require specific planning and management.

**Keywords:** Boosted Regression Tree, corridor, elephant habitat suitability, important value index, vegetation.

**Nepali Abstract:** नेपालमा पाइने जङ्गली हात्तीको वासस्थान र क्रियाकलाप विशेषतः चितवन राष्ट्रिय निकुञ्जको पूर्वी क्षेत्रमा पर्ने पर्सो राष्ट्रिय निकुञ्ज, यसको मध्यवर्ती क्षेत्रमा रहेको छ। यस अनुसन्धानले जङ्गली हात्तीको वासस्थानको उपयुक्तता, हात्तीले प्रयोग गर्ने क्षेत्र र हात्तीको उपस्थितिका कारकको बारेमा अध्ययन गरेको छ। स्थलगत रूपमा ६७ वटा प्लटको स्थापना गरी उक्त प्लटभित्र रहेका वनस्पतीहरूको विश्लेषण गरिएको थियो। बुस्टेड रिग्रेसन ट्री विधि लागू गरेर हात्तीको वासस्थान उपयुक्तता र कारकहरू जसमा स्तलाकृति (उचाइ, भिरालो, मोहडा), जलवायु (वर्षा, तापक्रम), वासस्थान प्राथमिकता, जमीन आवरण, छत्र आवरण जस्ता विभिन्न कारकहरूको अध्ययन गरिएको थियो। अध्ययनको परिणामले हात्तीको वासस्थान मुख्यतया प्रमुख वनस्पती प्रजाती, तापक्रम, उचाइ, वासस्थान प्राथमिकता र वर्षा निर्धारण गर्छ भनेर जनाएको छ। भिरालोपन, जमीन आवरण, छत्र आवरण र अरु कारकहरूले हात्तीको वासस्थान निर्धारणमा कम असर गर्ने देखियो। अध्ययन गरिएको क्षेत्रबाट ४०० मि. उत्तरपूर्वी र दक्षिणपूर्वी मोहडासम्म हात्ती सकृय रहेको पाइयो। न्युन भिरालोपन भएका खयर (*Acacia catechu*) र कालीकाठ (*Myrsine semicarpata*) प्रजातिले हात्तीको वनक्षेत्र जहाँ वार्षिक वर्षा ३०० मिमि रहन्छ त्यस्तो क्षेत्र सबैभन्दा उपयुक्त वासस्थानको रूपमा पाइएको छ। उल्लेखित क्षेत्र तोकेर यस मोडेलिङले हात्तीको वासस्थानको वातावरण उपयुक्तता र पर्सो र यसको मध्यवर्ती क्षेत्रमा हात्तीको संरक्षणको लागी ज्ञान प्रदान गर्नतर्फ जोड दिएको छ, जुन कुरा हात्तीको वासस्थान व्यवस्थापन र योजनामा आवश्यक पर्छ।

**DOI:** <https://doi.org/10.11609/jott.4467.11.13.14643-14654>

**Editor:** L.A.K. Singh, Bhuaneswar, Odisha, India

**Date of publication:** 26 October 2019 (online & print)

**Manuscript details:** #4467 | Received 02 August 2018 | Final received 27 June 2019 | Finally accepted 18 September 2019

**Citation:** Sharma, P., H. Adhikari, S. Tripathi, A.K. Ram & R. Bhattarai (2019). Habitat suitability modeling of Asian Elephant *Elephas maximus* (Mammalia: Proboscidea: Elephantidae) in Parsa National Park, Nepal and its buffer zone. *Journal of Threatened Taxa* 11(13): 14643–14654; <https://doi.org/10.11609/jott.4467.11.13.14643-14654>

**Copyright:** © Sharma et al. 2019. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by adequate credit to the author(s) and the source of publication.

**Funding:** None.

**Competing interests:** The authors declare no competing interests.

For **Author details** & **Author contribution** see end of this article.

**Acknowledgements:** Permission for the study was acquired from the Department of National Park and Wildlife Conservation, the government of Nepal. We would like to thank Hari Bhadracharya (Chief Conservation Officer, Parsa National Park (PNP) for his support during this study. This article is based on the bachelor thesis submitted to Agriculture and Forestry University, Faculty of Forestry, Hetauda, Nepal in partial fulfillment of the requirements for the degree of Bachelor of Forestry Science. Anonymous reviewers and the Journal Editor are thanked for useful insights and comments that helped improve this work.

## INTRODUCTION

The Asian Elephant was recognized as an endangered species in 1975 after its inclusion in Appendix I of CITES (Convention on International Trade in Endangered Species of Wild Fauna and Flora) (Bisht 2002) and listed as “Endangered” on the IUCN Red List of Threatened Species (IUCN 2017). These elephants are found in a variety of habitats that include grasslands, tropical evergreen, moist deciduous, dry deciduous and dry thorn forests, as well as secondary forest, scrublands, and cultivated areas (Sukumar 2003). Armbruster & Lande (1993) stated that human encroachment of natural habitats is one of the most critical issues facing elephant conservation. In Asia, elephants have lost extensive habitat areas, and as a result, conflicts with people have increased (Santiapillai 1997).

In Nepal, elephants are distributed throughout the lowland Terai in four isolated populations ranging over 10,982km<sup>2</sup> of forest habitat (DNPWC 2008). The estimated number of resident wild elephants in Nepal is between 107 and 145 (DNPWC 2008; Pradhan et al. 2011). The eastern population has 7–15 resident animals and 100 migratory animals from India. In central Nepal, 20–25 elephants reside primarily in Parsa National Park (PNP) and Chitwan National Park (CNP). The western and far western populations consist of 60–80, and 15–20, wild elephants respectively (DNPWC 2008; Pradhan et al. 2011).

Habitat conservation is an important aspect of wildlife conservation, and habitat suitability analysis is an essential aspect of management of wild animals such as elephants. Habitat suitability modeling can predict the quality and suitability of habitats for given species based on predictor variables such as topography (aspect, slope, altitude), climate (temperature, precipitation) and other biotic and abiotic factors. Different methods of modeling are used to determine suitable habitats for elephants. The boosted regression trees (BRT) method is an ensemble tree-based species distribution modelling technique that iteratively grows small/ simple trees based on the residuals from all previous trees (Elith et al. 2008). BRT has proven useful for working with large datasets of environmental variables and observations (Elith et al. 2008). For example it has been used to identify determinants of above ground biomass (Adhikari et al. 2017) and fish species distribution (Elith et al. 2008; Trigal & Degerman 2015). BRT and geographic information have also proven to be effective in the assessment of habitat quality.

The present study aims: 1) to assess the suitable

habitat of elephants, and 2) to determine which explanatory variables better explain elephant presence in PNP and buffer zone. This study has assessed habitat suitability in order to provide insights towards better management of elephant populations.

## METHODS AND MATERIALS

### Study area

The study was conducted in Parsa National Park (PNP) and its buffer zone (BZ), located in the sub-tropical zone of the southern part of Nepal. It has an area of 627km<sup>2</sup>. In 1984, PNP was established to preserve the habitat of natural populations of Asian Elephant *Elephas maximus*, Tiger *Panthera tigris*, and Gaur *Bos gaurus* (Rimal et al. 2018). The BZ of PNP was declared in 2005, which covers an area of 285.17km<sup>2</sup> encompassing three districts and 11 village development committees (VDC) (Figure 1). The region experiences four different seasons: summer (April–June), rainy/monsoon (July–September), winter (October–December), and spring (January–March).

The forests of PNP consist of tropical and subtropical tree species. Sal *Shorea robusta* forests compose about 90% of the park's vegetation. The riverine forests are found along the banks of rivers entailing species like Sisso *Dalbergia sisoo*, Silk Cotton Tree *Bombax ceiba*, and Khair *Acacia catechu*. Grass including Siru *Imperata cylindrica* and Kans *Saccharum spontaneum* are in the park. PNP and BZ support various endangered animal species including wild Asian Elephant, Royal Bengal Tiger *Panthera tigris*, and Sloth Bear *Melursus ursinus*. Mammals including Blue Bull *Boselaphus tragocamelus*, Sambar *Rusa unicolor*, Hog Deer *Axis porcinus*, Barking Deer *Indian muntjac*, Rhesus Macaque *Macaca mulatta*, and Palm Civet *Paradoxurus hermaphrodites* are also found in the park. Anthropogenic pressures like sand extraction, shifting cultivation and domestic cattle grazing are high in PNP and BZ (CHEC Nepal 2012).

### Quantitative data collection

Field work was conducted during the morning hours of May–June 2017. In reconnaissance, the habitats preferred by elephants were identified in consultation with local people. Questions concerned areas where elephants were frequently sighted, places where indirect signs of elephants were found, and availability of water. Reconnaissance field visits were made with the help of elephant rides, on foot and by vehicle, and areas were allocated into blocks according to habitat types. Sample plot centers were positioned using hand-held Garmin






Figure 1. Location of the study area and sample plot distribution in Parsa National Park and buffer zone, Nepal.

global positioning system (GPS) with a 2–5 m accuracy.

Nested quadrats of different size were purposively assigned in the study area (Figure 2). Total 67 plots were assigned and used to assess the status of tree, pole and regeneration condition. Quadrats of 10m × 10m were set in the study area to calculate the intensity for tree species. All plant species within each quadrat were identified and counted. For trees, trunk diameter at breast height (DBH; 1.3m) and height was measured. Quadrats of 5m × 5m were allocated randomly for shrubs. Herbs and regeneration were recorded from nesting sampling of 1m × 1m quadrat within the 5m × 5m quadrat.

Tree diameter, height, dominant species, crown cover and ground cover were measured, poles and regeneration were counted and in cases of grasses, clumps were counted within each quadrat. Plant species were identified by a local para-taxonomist, field guide and also based on literature related to plant identification in Nepal (Rimal et al. 2018). Leaves of unidentified tree species were brought to the faculty of forestry at Agriculture and Forestry University (AFU) and were identified.

To assess the habitat, important value index (IVI) and prominence value (PV) of vegetation available in the habitat range is crucial. The vegetation data collected in the field were used to calculate IVI, density, relative density, frequency, and relative frequency of the tree species by using equations 1–8 explained in Greig-Smith (1983). The IVI of a species signifies its dominance and




Figure 2. The layout of the quadrat used to assess the status of tree, pole and regeneration condition.

ecological success, its good power of regeneration and greater amplitude. The IVI was calculated by using three measures including relative frequency, relative density, and relative dominance. Vegetation data were calculated following the broad principle described by Mishra (1968) and Mueller-Dombois & Ellenberg (1974). Basal area helps to determine the dominance and nature of the community and it refers to the actual ground covered by the stems. Density is generally used for large plants that have discrete individuals (Zobel et al. 1987). Frequency and relative frequency give an index on the spatial distribution of a species (Krebs 1978). To calculate the

prominence value, the percentage cover of each species is estimated in each quadrat and recorded in classes as follows, for high coverage = > 50%, medium = 26–50% and low = 0–25%. Prominence value is used to calculate the availability of plants in the research sites (Jnawali 1995).

1. Density of species = (Total number of individuals of a species) / (Total number of quadrats sampled × area of a quadrat)

2. Relative density of species (RD) = (Total individuals of species) / (Total individual of all species)

3. Frequency of species = Number of plots in which a particular species occurs / Total number of plots sampled × 100

4. Relative frequency of species (RF) = Frequency of a species / Total frequency of all species × 100

5. Relative dominance of species = Total basal area of a species / Total basal area of all species × 100

6. Basal area =  $\pi d^2/4$

7. Important value index (IVI) = Relative density + Relative frequency + Relative dominance

8. PVX = MX (VFX)

(where, d = diameter at breast height (1.3 m) of tree, PVX = prominence value of species X; MX = mean percentage cover of species X; FX = Frequency of occurrence of species X)

#### Explanatory variables for modeling of habitat suitability distribution

A range of explanatory variables was derived from geospatial data sets for modeling habitat suitability. Table 1 presents the complete list of variables. The slope, aspects, and altitude were derived from the Japan Aerospace Exploration Agency (JAXA) digital elevation model (dem). Precipitation and temperature were downloaded from WorldClim data. Field measurements, dominant species, habitat preference, segment type, crown and ground cover and substrate conditions were derived. All topographic, climatic, and land use data available for the study area were resampled to 30m resolution and UTM 45N, WGS 84 projection system. For each absence and presence of GPS location, these variables were extracted. The correlation co-efficient between the explanatory variables and presence-absence data of elephant is shown in Appendix 1.

#### Statistical analysis

Boosted regression tree (BRT) (Elith et al. 2008) was used for examining the habitat preference area for elephant. BRT handles different types of predictor variables and accommodates missing data (Elith et al.

2008). Besides these, there is no need for prior data transformation or elimination of outliers. This is an advanced form of regression methods, which consists of two component—regression trees and boosting. BRT analysis was done using the ‘dismo’ package in R. The Bernoulli error distribution was used. Furthermore, the minimum predictive error was achieved when using a learning rate of 0.001, tree complexity (interaction depth) of 5, bag fraction of 0.75 and tolerance method “fixed”. All predictor variables were used as BRT can handle multi-collinearity among variables.

## RESULTS

#### Habitat assessment

IVI was calculated to find out the dominant tree species (Appendix 2) and prominence value was observed in the case of shrubs and herbs (Appendix 3 & 4). We calculated the species diversity of the study area for trees. Fifty-seven species were present in the quadrates; among them, 10 tree species were the dominant tree species present in the study area. In the study area, Sal (IVI=50.7753) was found most dominant and *Careya arborea* (IVI=5.2802) as least dominant. The species including *Mallotus philipinensis*, *Dillenia pentagaina*, and *Careya arborea* have the highest IVI among all, and they are the most preferred species of the elephant.

To determine the preferred habitat used by elephant, we calculated the PV of shrubs and herbs. Among 40 species, each of shrubs and herbs was present in the study area. Among the shrub species, *Eupatorium spp.* (PV=306.25) was the most abundant species and *Bauhinia vahilli* (PV=53.07) was the least abundant. Among the herb species, *Imperata cylindrica* (PV=317.66) was the most abundant and *Piper longum* (PV=29.48) the least abundant. The shrub species including *Eupatorium odoratum*, *Leea macrophylla*, and *Clerodendron viscosum* and herb species including *Imperata cylindrica*, *Saccharum spontaneum*, and *Fritillaria camschatcensis* have the highest PV among all species found in the study area as well as, they are the most preferred species of the elephant in the study area.

#### Habitat suitability

The total deviance explained by the BRT model was 0.16. The correlation between different variables, including presence-absence, altitude, land cover of the plot, segment type, substrate condition, dominant species, ground cover, crown cover, habitat preference,

**Table 1. Predictor variables used to model the habitat of Elephant.**

Predictor variables	Format (Source)	Description
Temperature ( $\times 10^\circ\text{C}$ ) (1km $\times$ 1km)	Raster (WorldClim) 1*	The temperature of June was used
Precipitation (mm) (1km $\times$ 1km)	Raster (WorldClim) 1*	Precipitation of June was used
Slope ( $^\circ$ ) (30m $\times$ 30m)	Raster (Jaxa DEM) 2*	
Aspect (30m $\times$ 30m)	Raster (Jaxa DEM) 2*	
Altitude (30m $\times$ 30m)	Raster (Jaxa DEM) 2*	
Habitat preference	Field measurement	Species preferred by elephant, including <i>Mallotus philippinensis</i> , <i>Imperata cylindrica</i> , <i>Dillenia pentagyna</i> , <i>Saccharum spontaneum</i> , <i>Careya arborea</i> , and <i>Pennisetum purpureum</i>
Dominant Species	Field measurement	Area dominated by species like <i>Acacia catechu</i> (AcC), <i>Bombax ceiba</i> (BoC), <i>Dillenia pentagyna</i> (DiP), <i>Albizia procera</i> (AlP), <i>Lagerstroemia parviflora</i> (LaP), <i>Terminalia chebula</i> (TeC), <i>Trewia nudifolia</i> (TrN), and <i>Myrsine semicerata</i> (MyS)
Segment type	Field measurement	Divides the area into the segment by fire line, foot trail, pond, river, and railway
Crown and ground cover	Field measurement	Cover (%) of forest crown and ground
Substrate condition	Field measurement	The condition of the soil, including hard soil, soft soil, and leaf litter

1\*—[www.worldclim.org](http://www.worldclim.org) | 2\*—[www.global.jaxa.jp/press/2015/05/20150518\\_daichi.html](http://www.global.jaxa.jp/press/2015/05/20150518_daichi.html).

precipitation, temperature, slope, and aspect is shown in Appendix 1. The relative influence of each predictor variables is shown in Figure 3. Each predictor variable has different relative contributions for the BRT model. Dominant species, temperature, and altitude have higher relative influence, whereas ground cover, crown cover and substrate condition have a lower relative influence.

Prioritized dependence plots visualize the effect of a single variable on the model response, holding all other variables constant. Model results vary the most with dominant species as seen in the first left plot (Figure 3). Dominant species (34%), temperature (15.3%), altitude (12.4%), habitat preference (11.1%) and precipitation (8.8 %) have the highest relative influence percentage and play a crucial role in the elephant distribution based on these plots. For more details on how they were calculated and model parameters used, see Sharma (2017).

On the basis of partial dependence plots, the elephant was more available at the altitude of 250–350 m with precipitation 310mm. The suitable habitat for the elephant was at the temperature of 28.5°C, a slope of 0–5° and in the northeastern and southeastern regions. Dominant species shows that *Acacia catechu* (AcC) and *Myrsine semicerata* (MyS) forest are more suitable for the elephant. Species including *Dillenia pentagyna* (DiP), *Saccharum spontaneum* (SaS), and *Pennisetum purpureum* (PeP) are the most preferable species of the elephant. Elephants dwelled in forest dominated by *Mallotus philippinensis* followed by *Syzygium cumini*. Thus areas having these species were the most suitable

habitats.

The weighted mean of discrete data was not available, whereas the weighted mean of continuous data was altitude (264m), precipitation (310mm), temperature (28.6 °C), slope (5.6°) and aspect (190°). Elephants were found mostly around the fire line and river, at an altitude of 150–350 m with temperature around 28.6 °C, crown cover 40–70 and slope below 0–5° (Figure 4).

The correlation between elephant presence-absence and temperature is 0.24, that implies a slight positive relationship between them, the elephant is mostly found in increasing temperature (Appendix 1). Whereas there is almost no linear association between presence-absence, and slope, dominant species, land cover of the plot, crown cover and ground cover. The relationship of altitude with temperature is negative, i.e., 0.84, the temperature of the area increases with a decrease in altitude and vice versa.

The elephant distribution prediction map based on altitude, slope, aspect, precipitation, and temperature only using boosted regression tree model is presented in Appendix 5. Other predictor variables were based specifically on field data and their extrapolation to spatial scale was not possible.

## DISCUSSION

The elephant population in Nepal is restricted to the Terai and Siwalik regions, where there have been large-scale conversion of forest and expansion of agricultural lands (Koirala et al. 2015). This has resulted






Figure 3. Single-variable partial dependence plots and smoothed response curves for the predictor variables, where Y-axis is the fitted function for the response variable (elephant). The relative influence is shown in parentheses.



in negative human-elephant interactions in many parts of Nepal. Movement of elephants outside the national park and wildlife reserve could have been the result of unsuitable habitat, reduced supply of food and water, and encroachment by human beings. Assessing habitat suitability of elephants assists in the preparation of sustainable management plans. PNP and BZ has been the habitat of elephant for long time, but habitat suitability studies are rare in this area. This research examines habitat suitability of the elephant in PNP and BZ based on different variables including dominant species, temperature, altitude, habitat preference, precipitation, segment type, aspect, slope, ground cover, crown cover and substrate condition.

Based on the BRT model, PNP and BZ are suitable habitats for elephants. We witnessed the outcome of parameters as per the physical, biological and climatic features of the area like slope, aspect, altitude, precipitation, temperature, habitat preference, crown cover and ground cover. The result shown by Koirala et al. (2016) posits that species like *Spatholobus parviflorus*, *Saccharum spontaneum*, *Shorea robusta*, *Mallotus philippensis*, *Garuga pinnata*, *Litsea mono-petala* contributed the highest proportion of diet for an elephant in the PNP. In consent with our result, similar species of trees, shrubs, and herbs have the highest IVI and PV and distributed in the lower part of the area. This result concludes that the study area is the most suitable for elephant to dwell. Our study revealed that the habitat is suitable in the Northeast and Southeast region of the study area, which is similar to the result of Shamsuddoha & Aziz (2015).

Rood et al. (2010) studies have found that the elephant's habitat use in a tropical forest is depicted by areas of high forest cover. Our analysis, however, found no marked relationship between ground cover, crown cover, and presence of elephants. Our findings revealed that in the study area, slope 0–5° and altitude 400m is suitable for elephants which are almost similar to the result of Areendran et al. (2011). In accordance with the studies of Douglas et al. (2006), Lin et al. (2008) and Ochieng (2015), suitable habitat for elephants was found to be limited by augmentation of both altitude and slope. There is no abundance of elephants' presence sign with the increment of the altitude and slope in PNP. In order to preserve their energy needs, Ntumii et al. (2005) mention that elephants avoid the height and steeply sloped area.

Variability in results might have occurred due to the differences in sampling methods, variance in forest condition, composition, and sampling area, etc. The

research outcome was concluded based on only one season field work; however, taking all the results of four seasons might produce more effective result. Data of precipitation and temperature were extracted from Worldclim; the data taken from the nearest metrological station of Samara could be better with more accuracy. The outcomes from this study, linked to slope, and elevation are valid for PNP only, and cannot be generalized to the habitat of an elephant in other countries. Further research should focus on creating map of elephant distribution, habitat suitability, and threats to elephant from invasive species.

## CONCLUSION

BRT was applied to assess elephant habitat suitability in PNP. In this study, we analyzed the distribution of elephant using a combination of biotic and abiotic environmental variables, including the topographic and climatic factors. The model emphasizes on environmental suitability and contributes to knowledge for conservation of elephant in PNP. It provides a basis for habitat analysis. Elephants were recorded up to 400m and in northeastern and southeastern aspects. Its presence could not be related to forest cover and substrate condition. The result from the modeling may become useful to plan and delineate areas for management of elephant. It presents scope to minimize HEC through precautionary measures.


## REFERENCES

- Adhikari, H., J. Heiskanen, M. Siljander, E. Maeda, V. Heikinheimo & P.K.E. Pelikka (2017). Determinants of aboveground biomass across an Afromontane landscape mosaic in Kenya. *Remote Sensing* 9(8): 827. <https://doi.org/10.3390/rs9080827>
- Areendran, G., K. Raj, S. Mazumdar, M. Munsri, H. Govil & P.K. Sen (2011). Geospatial modelling to assess elephant habitat suitability and corridors in northern Chhattisgarh, India. *Tropical Ecology* 52: 275–283.
- Armbruster, P. & R. Lande (1993). A population viability analysis for African Elephant (*Loxodonta africana*): how big should reserves be? *Conservation Biology* 7(3): 602–610. <https://doi.org/10.1046/j.1523-1739.1993.07030602.x>
- Bisht, S.S. (2002). An overview of elephant conservation in India. *Indian Forester* 128(2): 121–136
- CHEC Nepal (2012). An assessment of Tiger (*Panthera tigris*) and its habitat in Bara forest, Nepal. A report submitted to District Forest Office, Simara, Bara.
- DNPWC (2008). The Elephant Conservation Action Plan for Nepal. Ministry of Forest and Soil Conservation, Department of National Parks and Wildlife Conservation, Kathmandu, Nepal, 30pp.
- Douglas-Hamilton, I., S. Bhalla, G. Wittemyer & F. Vollrath (2006). Behavioural reactions of elephants towards a dying and deceased matriarch. *Applied Animal Behaviour Science* 100: 87–102. <https://doi.org/10.1016/j.applanim.2006.04.014>

- Elith, J., J.R. Leathwick & T. Hastie (2008). A working guide to boosted regression trees, (Ml), 802–813. <https://doi.org/10.1111/j.1365-2656.2008.01390.x>
- Greig-Smith (1983). *Quantitative Plant Ecology*. University of California Press, Berkeley, CA, xiv+359pp.
- IUCN (2017). The IUCN Red List of Threatened Species. Version 2017-1. <https://www.iucnredlist.org>
- Jnawali, S.R. (1995). Population ecology of greater one horned rhinoceros (*Rhinoceros unicornis*) with particular emphasis on habitat preference, food ecology and ranging behavior of a reintroduced population in Royal Bardia National Park in Low land Nepal (A Doctor Scientiarum Thesis 1995: 4), i-vii, 128pp.
- Koirala, R.K., W. Ji, A. Aryal, Jessica Rothman & D. Raubenheimer (2015). Dispersal and ranging patterns of the Asian Elephant (*Elephas maximus*) in relation to their interactions with humans in Nepal. *Ethology Ecology & Evaluation* 28(2): 221–231. <https://doi.org/10.1080/03949370.2015.1066872>
- Koirala, R.K., D. Raubenheimer, A. Aryal, M.L. Pathak & W. Ji (2016). Feeding preferences of the Asian Elephant (*Elephas maximus*) in Nepal. *BMC Ecol.*, 16 (1): 54. <https://doi.org/10.1186/s12898-016-0105-9>
- Krebs, C.J. (1978). *Ecology: The Experimental Analysis of Distribution and Abundance*. 2<sup>nd</sup> Ed. Harper & Row Publishers, 678pp.
- Lin, L., L. Feng, W. Pan, X. Guo, J. Zhao, A. Luo & L. Zhang (2008). Habitat selection and the change in distribution of Asian elephants in Mengyang Protected Area, Yunnan, China. *Acta Theriologica* 53: 365–374. <https://doi.org/10.1007/BF03195197>
- Mishra, R. (1968). *Ecology Work Book*. Oxford and IBH Publishing Company, Calcutta, 244pp.
- Mueller-Dombois, D. & H. Ellenberg (1974). *Aims and Methods of Vegetation Ecology*. John Wiley & Sons, New York, 547pp.
- Ntumi, C.P., R.J. van Aarde, N. Fairall & W.F. de Boer (2005). Use of space and habitat use by elephants (*Loxodonta africana*) in the Maputo Elephant Reserve, Mozambique. *South African Journal of Wildlife Research* 35(2): 139–146.
- Ochieng, E.O. (2015). Characterizing the spatial distributions of elephants in Mpala, Kenya. MSc Thesis, University of Twente, 53pp.
- Pradhan, N.M.B., A.C. Williams & M. Dhakal (2011). Current status of Asian elephants in Nepal. *Gajah* 35: 87–92.
- Ram, A.K. (2008). Impact of *Mikania micrantha* on *Rhinoceros unicornis* habitat in Chitwan National Park, Chitwan Nepal (Bachelor thesis submitted to Tribhuvan University), 79pp. <https://doi.org/10.13140/RG.2.1.1756.8244>
- Rimal, S., H. Adhikari & S. Tripathi (2018). Habitat suitability and threat analysis of Greater One-horned Rhinoceros *Rhinoceros unicornis* Linnaeus, 1758 (Mammalia: Perissodactyla: Rhinocerotidae) in Rautahat District, Nepal. *Journal of Threatened Taxa* 10(8): 11999–12007. <https://doi.org/10.11609/jott.3948.10.8.11999-12007>
- Rood, E., A.G. Abdullah & V. Nijman (2010). Using presence-only modelling to predict Asian Elephant habitat use in a tropical forest landscape: implications for conservation. *Diversity and Distributions* 16: 975–984. <https://doi.org/10.1111/j.1472-4642.2010.00704.x>
- Santiapillai, C. (1997). The Asian Elephant conservation: a global strategy. *Gajah* 18: 21–39.
- Shamsuddoha, M. & M.A. Aziz (2017). Raiding pattern of migratory elephants in a human dominated landscape in northern Bangladesh. *Ecoprint* 24: 21–27. <https://doi.org/10.3126/eco.v24i0.20643>
- Sharma, P. (2017). Habitat suitability modeling of Asian wild elephant and its interaction with people in Parsa National Park and its Buffer Zone. Submitted to Agriculture Forest University, Faculty of Forestry, Hetauda, Nepal in partial fulfilment of the requirement for the degree of Bachelors' in forestry, i-xi, 46pp.
- Sukumar, R. (2003). *The living elephants: evolutionary ecology, behavior, and conservation*. Oxford University Press, New York, 478pp.
- Trigal, C. & E. Degerman (2015). Multiple factors and thresholds explaining fish species distributions in lowlands streams. *Global Ecology and Conservation* 4: 589–601. <https://doi.org/10.1016/j.gecco.2015.10.009>
- Zobel, D.B., U.K. Yadav, P.K. Jha & M.J. Behan (1987). *A practical manual for ecology*. Rani Printing Press, Kathmandu, Nepal.

**Author details:** PUJA SHARMA is doing MSc in Tropical and International Forestry at University of Gottingen, Germany. She has completed BSc in forestry from Agriculture and Forestry University and worked in the field of Natural Resource Management and wildlife. HARI ADHIKARI defended his PhD at University of Helsinki, Finland. He has international working experience on wildlife and forestry in Nepal, India, Philippines, Kenya, Germany and Finland. SHANKAR TRIPATHI is young faculty member of Faculty of Forestry at AFU, who have involve in lecturing and research on forest measurement and application of Remote sensing and GIS in the field of nature conservation. He has accomplished post graduate degree in Forestry from Tribhuvan University. ASHOK KUMAR RAM holds MSc on Forestry from Institute of Forestry Pokhara, Nepal and studying his PhD on Wildlife Science from Wildlife Institute of India. He is an IUCN Species survival commission members for Asian Elephant specialist group. RAJEEV BHATTARAI is a graduate student in School of Forest Resources, University of Maine with a undergraduate degree in forestry from Tribhuvan University, Nepal.

**Author contribution:** PS planned and conducted this research, HA and ST supervised this research. HA, PS and ST together worked on manuscript. HA and PS collected RS and GIS data. PS, RB and ST collected field data. AKR supervised PS during field data collection.



Appendix 1. Pearson correlation between different predictor variables and their scatter plots.



**Appendix 2. Important Value Index (IVI) of dominance tree species**


Scientific name	Local name	Relative frequency	Relative density	Relative dominance %	Important value index (IVI)
<i>Shorea robusta</i>	Sal	13.36	0.27	37.14	50.77
<i>Mallotus philipinensis</i>	Sindure	6.45	0.14	10.60	17.20
<i>Terminalia tomentosa</i>	Asna	4.60	0.04	8.22	12.87
<i>Acacia catechu</i>	Khair	4.60	0.11	5.57	10.29
<i>Lagestomia parviflora</i>	Bot dhayero	6.91	0.05	3.44	10.41
<i>Dillenia pentagyna</i>	Tatari	6.45	0.05	3.73	10.23
<i>Adina cordifolia</i>	Haldu	3.68	0.02	3.44	7.14
<i>Garuga pinnata</i>	Dabdbe	5.06	0.03	1.97	7.08
<i>Albigia procera</i>	Setosiris	2.76	0.01	3.10	5.88
<i>Careya arborea</i>	Kumbe	2.76	0.01	2.50	5.28

**Appendix 3. Prominence value of shrubs in the study area.**

Scientific name	Local name	Number of individuals	Frequency	Mean cover of individual species ( $M_x$ )	Prominence value ( $PV_x$ )
<i>Eupatorium odoratum</i>	Setobanmara	188	50.72	43	306.25
<i>Leea macrophylla</i>	Galini	213	50.72	38	270.64
<i>Clerodendron viscosum</i>	Bhati	248	33.33	25	144.33
<i>Murraya koenigii</i>	Curry leaf	72	26.08	20	102.15
<i>Fritillaria</i> spp.	Thulobandhan	19	5.79	30	72.23
<i>Lantana camara</i>	Lantana	14	4.34	30	62.55
<i>Asparagus racemosus</i>	Kurilo	34	18.84	14	60.76
<i>Agiratus conyzoides</i>	Gande	111	5.79	25	60.19
<i>Parthenium</i> spp.	Parthenium	517	5.79	24	57.78
<i>Bauhinia vahlii</i>	Bhorla	20	8.69	18	53.07

**Appendix 4. Prominence value of herbs in the study area.**

Scientific name	Local name	Number of individuals	Frequency	Mean cover of individual species ( $M_x$ )	Prominence value ( $PV_x$ )
<i>Imperata cylindrica</i>	Siru	3516	42.02	49	<b>317.66</b>
<i>Saccharum spontaneum</i>	Kans	970	13.04	32	115.57
<i>Fritillaria camschatcensis</i>	Ban dhan	1035	18.84	20	86.81
<i>Hemaltriya compressa</i>	Ghodaydubo	699	10.14	25	79.62
<i>Cynodon dactylon</i>	Dubo	634	7.24	29	78.06
<i>Digitaria</i> spp.	Chitrebanso	382	10.14	23	73.25
<i>Pennisetum purpureum</i>	Elephant grass	163	4.34	22	45.87
<i>Barlaria cristata</i>	Kuro	151	7.24	16	43.07
<i>Dendrobium</i> spp.	Orchid	27	7.24	11	29.61
<i>Piper longum</i>	Pipla	18	8.69	10	<b>29.48</b>



Appendix 5. Elephant distribution prediction based on altitude, slope, aspect, precipitation, and temperature only using boosted regression tree model. Other predictor variables were based on field data and not available at the wall-to-wall spatial scale.







www.threatenedtaxa.org



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

October 2019 | Vol. 11 | No. 13 | Pages: 14631–14786

Date of Publication: 26 October 2019 (Online & Print)

DOI: 10.11609/jott.2019.11.13.14631-14786

## Communications

### Camera trap survey of mammals in Cleopatra's Needle Critical Habitat in Puerto Princesa City, Palawan, Philippines

– Paris N. Marler, Solomon Calago, Mélanie Ragon & Lyca Sandra G. Castro, Pp. 14631–14642

### Habitat suitability modeling of Asian Elephant *Elephas maximus* (Mammalia: Proboscidea: Elephantidae) in Parsa National Park, Nepal and its buffer zone

– Puja Sharma, Hari Adhikari, Shankar Tripathi, Ashok Kumar Ram & Rajeev Bhattarai, Pp. 14643–14654

### Current population status of the endangered Hog Deer *Axis porcinus* (Mammalia: Cetartiodactyla: Cervidae) in the Terai grasslands: a study following political unrest in Manas National Park, India

– Alolika Sinha, Bibhuti Prasad Lahkar & Syed Ainul Hussain, Pp. 14655–14662

### A food spectrum analysis of three bufonid species (Anura: Bufonidae) from Uttarakhand region of the western Himalaya, India

– Vivekanand Bahuguna, Ashish Kumar Chowdhary, Shurveer Singh, Gaurav Bhatt, Siddhant Bhardwaj, Nikita Lohani & Satyanand Bahuguna, Pp. 14663–14671

### Moulting pattern and mortality during the final emergence of the Coromandel Marsh Dart Damselfly *Ceriatrion coromandelianum* (Zygoptera: Coenagrionidae) in central India

– Nilesh R. Thaokar, Payal R. Verma & Raymond J. Andrew, Pp. 14672–14680

### Diversity of parasitic Hymenoptera in three rice-growing tracts of Tamil Nadu, India

– Johnson Alfred Daniel & Kunchithapatham Ramaraju, Pp. 14681–14690

### Mapping octocoral (Anthozoa: Octocorallia) research in Asia, with particular reference to the Indian subcontinent: trends, challenges, and opportunities

– Ghosh Ramvilas, Kannan Shalu, Rajeev Raghavan & Kutty Ranjeet, Pp. 14691–14721

### SEM study of planktonic chlorophytes from the aquatic habitat of the Indian Sundarbans and their conservation status

– Gour Gopal Satpati & Ruma Pal, Pp. 14722–14744

### Is cultivation of *Saussurea costus* (Asterales: Asteraceae) sustaining its conservation?

– Chandra Prakash Kuniyal, Joel Thomas Heinen, Bir Singh Negi & Jagdish Chandra Kaim, Pp. 14745–14752

## Short Communications

### A first photographic record of a Yellow-bellied Weasel *Mustela kathiah* Hodgson, 1835 (Mammalia: Carnivora: Mustelidae) from western Nepal

– Badri Baral, Anju Pokharel, Dipak Raj Basnet, Ganesh Bahadur Magar & Karan Bahadur Shah, Pp. 14753–14756

### Mammal diversity in a montane forest in central Bhutan

– Tashi Dhendup, Kinga Thinley & Ugyen Tenzin, Pp. 14757–14763

## Notes

### First record of Otter Civet *Cynogale bennettii* (Mammalia: Carnivora: Viverridae) kept as a pet in Indonesia, representing a possible new threat to the species

– Jamie Francis Bernard Bouhuys, Pp. 14764–14766

### An observation of the White-bellied Sea Eagle *Haliaeetus leucogaster* preying on Saltwater Crocodile hatchlings *Crocodylus porosus* in Bhitarkanika Wildlife Sanctuary, India

– Nimain Charan Palei, Bhakta Padarbinda Rath & Bimal Prasanna Acharya, Pp. 14767–14769

### Elusive, rare and soft: a new site record of Leith's Softshell Turtle *Nilssonia leithii* (Reptilia: Testudines: Trionychidae) from

Bhadra Tiger Reserve, Karnataka, India  
– H.S. Sathya Chandra Sagar, M. Mrunmayee, I.N. Chethan, Manish Kumar & D.V. Girish, Pp. 14770–14772

### A new distribution record of the Pentagonal Sea Urchin Crab *Echinoecus pentagonus* (A. Milne-Edwards, 1879) (Decapoda: Brachyura: Pilumnidae) from the Andaman Islands, India

– Balakrishna Meher & Ganesh Thiruchitrabalam, Pp. 14773–14776

### First records of the ghost moth genus *Palpifer* Hampson, [1893] (Lepidoptera: Hepialidae) from the Indian subcontinent south of the Himalaya

– Siyad A. Karim & John R. Grehan, Pp. 14777–14779

### First record of longhorn beetle *Calothyrsa margaritifera* (Cerambycidae: Lamiinae: Phrynetini) from western India

– Vishwas Deshpande & Hemant V. Ghatge, Pp. 14780–14783

### Extended distribution of *Ceropegia mahabalei* Hemadri & Ansari (Apocynaceae) to the state of Gujarat, India

– Mukta Rajaram Bhamare, Hemantkumar Atmaram Thakur & Sharad Suresh Kambale, Pp. 14784–14786

## Partner



## Member



## Publisher & Host

